

Departamento de
SALUD



Guías de Servicios Pediátricos Preventivos

Revisada 2024

Aprobada por

Carlos R. Mellado Lopez M.D.
Secretario

Departamento de Salud

6 de agosto de 2024

Guías de Servicios Preventivos Pediátricos 2024

El Departamento de Salud establece las Guías de Servicios Pediátricos Preventivos con el propósito de promover las mejores prácticas al brindar servicios médicos a la población pediátrica. Las guías proveen recomendaciones para que toda persona de 21 años o menos reciba evaluaciones médicas que provean la oportunidad para identificar y diagnosticar tempranamente aquellas condiciones físicas, mentales y conductuales que requieren una pronta atención y en cumplimiento con los requisitos de EPSDT del Centro de Servicios para Medicaid. El contenido de las guías se desarrolló con recomendaciones por un panel de expertos, recomendaciones actualizadas de *Bright Futures Guidelines for Health Supervision of Infants, Children and Adolescents*, publicado por la Academia Americana de Pediatría (versión marzo 2021), y de *United States Preventive Task Force*.

Interpretación de Guías de Servicios Pediátricos Preventivos

Las Guías de Servicios Pediátricos Preventivos están divididas en cuatro etapas:

- Infancia de 0 a 9 meses
- Niñez Temprana de 12 a 48 meses
- Niñez de 5 a 10 años
- Jóvenes de 11 a 21 años

En cada etapa el esquema incluye las edades a las cuales se debe ofrecer una visita preventiva, y un listado de evaluaciones, cernimientos y/o pruebas de laboratorios universales, las cuales se recomiendan se administren a toda la población pediátrica en las edades indicadas por el símbolo En el listado se enfatizan áreas específicas a explorar en el historial y examen físico, temas a enfatizar durante la orientación anticipatoria, cernimientos recomendados a utilizar para completar la evaluación y acciones a tomar relevantes a los resultados, durante la visita médica.

Cada etapa presenta un listado de evaluaciones, cernimientos y/o pruebas de laboratorios selectivas, pruebas o acciones a tomar por el profesional de la salud justificado por el juicio clínico y los hallazgos de riesgo en las evaluaciones (historial, examen físico y/o cernimientos) indicando las edades correspondientes por el símbolo

Guías de Servicios Preventivos Pediátricos 2024

Universal Infancia	Acción	Visita Prenatal	Recién Nacido	3 a 5 días	1 mes	2 m	4 m	6 m	9 m
Historial / Inicial o intervalo	En visita prenatal historial familiar y de embarazo	✓	✓	✓	✓	✓	✓	✓	✓
Examen físico	En cada visita examen físico esencial		✓	✓	✓	✓	✓	✓	✓
Medidas: peso (kg), largo (cm), Circunferencia de cabeza (cm)	Clasificar y evaluar percentil en gráfica		✓	✓	✓	✓	✓	✓	✓
Vigilancia, desarrollo y evaluación de conducta y condición psicosocial	Observación clínica e historial, atención a los determinantes sociales, trauma, seguridad alimentaria		✓	✓	✓	✓	✓	✓	✓
Inmunizaciones	Evaluar cumplimiento con esquema vigente y administrar vacunas necesarias para su cumplimiento		✓	✓	✓	✓	✓	✓	✓
Evaluación y apoyo de lactancia/ orientación alimentación en 1er año	Vigilar aumento de peso en primera semana y referir a grupos de apoyo en la comunidad cuando sea indicado, orientar introducción de sólidos y alimentos	✓	✓	✓	✓	✓	✓	✓	✓
Cernimiento de Depresión Materna	Cernimiento Edinburgh/ Referir para apoyo y ayuda si resulta positivo					✓		✓	
Guía Anticipatoria	Enfatizar prácticas de dormir seguro y prevención de lesiones no intencionales	✓	✓	✓	✓	✓	✓	✓	✓
Cernimiento Auditivo Ley 311, 2003	A infantes con pruebas positivas deben realizárseles la prueba confirmatoria antes de los 3 meses de edad y recibir tratamiento definitivo antes de cumplir 6 meses de edad		✓						
Cernimiento Metabólico y Hemoglobinopatía	Ley 84, 1987		✓						
Cernimiento Defecto Cardíaco Congénito Crítico	Oximetría de pulso luego de las 24 horas de nacido, antes del alta		✓						
Cernimiento Hiperbilirrubinemia	Prueba de bilirrubina antes del alta de hospital, a las 48 horas nacidos en el hogar		✓						
Cernimiento del Desarrollo	Administrar instrumento de cernimiento validado. Ages and Stages (ASQ) última edición o Survey Wellbeing Young Children (SWYC)								✓
Evaluación y Cernimiento de riesgo de caries	Cernimiento de riesgo para caries (Caries-risk Assessment Questionnaire), resultado de alto riesgo, referir inmediatamente con el primer diente al dentista							✓	✓
Cernimiento Riesgo Tuberculosis	Cuestionario de riesgos, historial de exposición			✓				✓	
Cernimiento riesgo de exposición a plomo	Cuestionario de riesgos con resultado positivo, ordenar muestra de plomo en sangre							✓	✓

Infancia (continuación)

Selectivo Infancia	Evaluación	Acción	Visita Prenatal	Recién Nacido	3 a 5 días	1 mes	2m	4m	6m	9m
Presión Arterial	Historial positivo de riesgo	Presión arterial		<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
Visión	Historial y/o físico positivo	Oftalmólogo		<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
Anemia	Historial de prematurez	Hematocrito o Hgb						<input checked="" type="checkbox"/>		

Guías de Servicios Preventivos Pediátricos 2024

Universal Niñez Temprana	Acción	12 m	15 m	18 m	24 m	30 m	36 m	48 m
Historial / Inicial o intervalo	Historial de actividad física y alimentación	✓	✓	✓	✓	✓	✓	✓
Examen Físico	En cada visita examen físico esencial	✓	✓	✓	✓	✓	✓	✓
Medida peso/estatura	Clasificar percentil en gráfica	✓	✓	✓	✓	✓	✓	✓
Circunferencia de cabeza	Clasificar percentil en gráfica	✓	✓	✓	✓			
Guía Anticipatoria	Controlar tiempo exposición a consolas digitales y tv	✓	✓	✓	✓	✓	✓	✓
Vigilancia desarrollo y evaluación de conducta y condición psicosocial	Observación clínica e historial	✓	✓	✓	✓	✓	✓	✓
IMC / BMI	Clasificar percentil en gráfica				✓	✓	✓	✓
Presión Arterial							✓	✓
Inmunizaciones	Evaluar cumplimiento con esquema vigente y administrar vacunas necesarias para su cumplimiento	✓	✓	✓	✓	✓	✓	✓
Agudeza Visual	Evaluar objetivamente agudeza visual (ej.: cartilla Snellen)						✓	✓
Cernimiento auditivo	Audiometría							✓
Anemia	Hematocrito o Hgb	✓						
Autismo	Administrar instrumento <i>Modified Checklist for Autism in Toddlers</i> , (M-CHAT) o versión revisada (M-CHAT-R/F), seguir protocolo para autismo			✓	✓			
Cernimiento en el Desarrollo	Administrar instrumento de cernimiento validado. <i>Ages and Stages</i> (ASQ) última edición o <i>Survey Wellbeing Young Children</i> (SWYC)			✓	✓	✓		
Evaluación de Salud Oral	Referir al dentista para: limpieza cada 6 meses y barniz de Fluoruro	✓		✓	✓	✓	✓	✓
Cernimiento Riesgo Tuberculosis	Cuestionario de riesgos, historial positivo de exposición ordenar PPD	✓			✓		✓	✓
Plomo	Nivel de plomo en sangre	✓			✓			

Niñez Temprana (Continuación)

Selectivo Niñez Temprana	Evaluación	Acción	12 m	15 m	18 m	24 m	30 m	36 m	48 m
Presión Sanguínea	Historial positivo de riesgos	BP	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>		
Visión	Historial y/o físico positivo	Oftalmólogo	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>		
Audición	Historial y/o físico positivo	Audiólogo	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	
Anemia	Historial y/o físico positivo	Hematocrito		<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>		<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
Dislipidemia	Historial familiar de riesgo enfermedades cardiovasculares y físico positivo (obeso)	Panel lípidos en ayuna				<input checked="" type="checkbox"/>			<input checked="" type="checkbox"/>
Plomo *	Cernimiento de riesgo (Cuestionario de riesgo)	Niveles plomo en sangre, cuando cuestionario es positivo para riesgo de exposición						<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>

Guías de Servicios Preventivos Pediátricos 2024

Universal Niñez	Acción		5 años	6 años	7 años	8 años	9 años	10 años
Historial / Inicial o intervalo	Historial de actividad física y bienestar emocional		✓	✓	✓	✓	✓	✓
Examen Físico	En cada visita examen físico esencial		✓	✓	✓	✓	✓	✓
Peso, estatura, IMC / BMI	Clasificar percentil en gráfica		✓	✓	✓	✓	✓	✓
Presión arterial			✓	✓	✓	✓	✓	✓
Guía anticipatoria	Promover actividad física más de una hora diaria / disminuir tiempo en consolas a menos de 2 horas diarias		✓	✓	✓	✓	✓	✓
Vigilancia desarrollo	Observación clínica e historial		✓	✓	✓	✓	✓	✓
Vigilancia conducta	Observación clínica e historial		✓	✓	✓	✓	✓	✓
inmunizaciones	Evaluar cumplimiento con esquema vigente Administrar vacunas necesarias para cumplir con esquema		✓	✓	✓	✓	✓	✓
Agudeza visual	Evaluar objetivamente agudeza visual (ej.: cartilla Snellen)		✓	✓		✓		✓
Cernimiento auditivos	Audiometría		✓	✓		✓		✓
Dislipidemia	Cernimiento: panel lípidos en ayuna						Una vez entre 9 a 10 años	
Salud oral	Visita al dentista para limpieza y evaluación 2 veces al año		✓	✓	✓	✓	✓	✓
Cernimiento riesgo Tuberculosis	Cuestionario de riesgos, historial de exposición positiva ordenar PPD o prueba IGRA en sangre (solo a mayores de 4 años)		✓	✓	✓	✓	✓	✓
Selectivo	Evaluación	Acción	5 años	6 años	7 años	8 años	9 años	10 años
Visión	Historial y/o físico positivo	Referido oftalmólogo	☑	☑	☑	☑	☑	☑
Audición	Historial positivo	Referido Audiólogo	☑	☑	☑	☑	☑	☑
Anemia	Historial y/o físico positivo	Hematocrito o Hgb.	☑	☑	☑	☑	☑	☑
Plomo	Historial de riesgo a exposición a plomo	Niveles plomo en sangre	☑	☑				
Dislipidemia	Historial familiar de riesgo enfermedades cardiovasculares y físico positivo (obeso)	Panel lípidos en ayuna	☑	☑	☑			

Guías de Servicios Preventivos Pediátricos 2024

Universal Jóvenes	Acción	11-14 años	15-17 años	18-21 años
Historial detallado y examen físico	Historial de actividad física y bienestar emocional. Desarrollo características sexuales secundarias, menarquia, sueños mojados, historial actividad sexual, hábitos de dormir, acoso <i>bullying</i>)	Anual	Anual	Anual
Peso, estatura, IMC/BMI	Clasificar percentil en gráfica	Anual	Anual	Anual
Presión arterial		Anual	Anual	Anual
Vigilancia desarrollo	Observación clínica e historial	Anual	Anual	Anual
Evaluación conducta	Cernimiento conducta de riesgo, conocida como CRAFFT versión 2.1+N en inglés o en español	Anual	Anual	Anual
Evaluación presencia de Violencia y/o Depresión	Cernimiento de depresión, "Patient Health Questionnaire 9" (PHQ9), historial de violencia o agresión	Anual	Anual	Anual
Promoción estilos de vida saludables	(Alimentación, Actividad Física, actividad sexual responsable y saludable, prevención del uso de alcohol y sustancias controladas, prevención de fumar y cigarrillos electrónicos)	Anual	Anual	Anual
Guía anticipatoria	Anticipar cambios fisiológicos y emocionales típicos para cada etapa	Anual	Anual	Anual
inmunizaciones	Evaluar cumplimiento con esquema vigente. Administrar vacunas necesarias para cumplir con esquema	Anual	Anual	Anual
Visión	Evaluación objetiva utilizando tabla optométrica (ej.: cartilla Snellen)	Una vez entre los 11 a 14 años	Una vez entre los 15 a 17 años	Una vez entre los 18 a 21 años
Cernimiento auditivo	Cernimiento por audiometría que incluya alta frecuencia entre 6,000 a 8,000 Hz.	Una vez entre 11 a 14 años	Una vez entre los 15 a 17 años	Una vez entre los 18 a 21 años
Dislipidemia	Panel lípidos en ayuna	una vez entre los 9 a 11 años		Una vez entre los 17 a 21 años
Salud Oral	Visita al dentista para evaluación y limpieza profesional 2 veces al año	Anual	Anual	Anual
VIH, Ley 45 de 2016	Prueba de laboratorio con consentimiento previo e información del significado de los resultados positivo o negativo / a partir de los 13 años, repetir prueba cada 5 años	13 años	Una vez entre los 15 a 17 años	
Prueba en orina para Clamidia, Gonorrea	Universal NAAT en orina		Una vez entre los 15 a 17 años	
Sífilis (VDRL)	Laboratorio			Una vez entre los 18 a 21 años
Displasia cervical	Pap smear			A los 21 años

Guías de Servicios Preventivos Pediátricos 2024

Universal Jóvenes	Acción	11-14 años	15-17 años	18-21 años
Tuberculosis	PPD o IGRA en sangre			Una vez entre los 18 a 21 años
Hepatitis C	Muestra de sangre para presencia de anticuerpos a Hepatitis C			Una vez en después de los 17 años

Selectivo	Evaluación	Acción	Temprana: 11-14 años	Media: 15-17 años	Tardía: 18-21 años
Visión	Historial, físico, cernimiento positivo	Referir al Oftalmólogo	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
Audición	Historial, físico, cernimiento positivo	Referir al Audiólogo	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
Anemia	Historial y/o físico positivo	Hematocrito o Hgb	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
Embarazo	Sospecha embarazo	Prueba serológica	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
Tuberculosis	Exposición casos TB positivo	PPD o IGRA en sangre	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	
Prueba en orina para Clamidia, Gonorrea	Historial y/o físico positivo	Laboratorios NAATS	<input checked="" type="checkbox"/>		<input checked="" type="checkbox"/>
Sífilis (VDRL)	Historial y físico positivo	Laboratorio VDRL	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	

Enlaces de Referencia:

Enlaces para pruebas de cernimiento recomendadas

- MCHAT y MCHAT-R/F <https://mchatscreen.com/>
- Ages and Stages (ASQ) <http://agesandstages.com/>
- SWYC <https://www.floatinghospital.org/The-Survey-of-Wellbeing-of-Young-Children/Age-Specific-Forms>
- CRAFFT o Carlos <https://craftt.org/get-the-craftt/>

Cuestionario para Evaluar Riesgo de Tuberculosis en Poblaciones Pediátricas

- <http://www.salud.eov.pr/Dept-de-Salud/Documents/Cuestionario%20para%20Evaluar%20Riesgo%20de%20Tuberculosis%20en%20Poblaciones%20Pedi%C3%A1tricas.pdf>

Cuestionario para evaluar riesgo de plomo

- Cernimiento de depresión PHQ-9 para adolescentes
[https://aidsetc.org/sites/default/files/resources_files/PHQ-A%20Spanish II%20.pdf](https://aidsetc.org/sites/default/files/resources_files/PHQ-A%20Spanish%20II%20.pdf)

Otros cernimientos para desórdenes de salud mental

- <https://www.hiv.uw.edu/page/mental-health-screening/ihsd>

Herramienta para identificar otros cernimientos

- <https://screeningtime.org/star-center/#/screening-tools>

Enlaces para herramientas en las visitas preventivas

Manual de AAP, Códigos para facturar visitas pediátricas

<https://downloads.aap.org/AAP/PDF/Coding%20Preventive%20Care.pdf>

Bright Futures AAP

- <https://brightfutures.aap.org/Pages/default.aspx>
- https://downloads.aap.org/AAP/PDF/periodicity_schedule.pdf

Each child and family is unique; therefore, these Recommendations for Preventive Pediatric Health Care are designed for the care of children who are receiving nurturing parenting, have no manifestations of any important health problems, and are growing and developing in a satisfactory fashion. Developmental, psychosocial, and chronic disease issues for children and adolescents may require more frequent counseling and treatment visits separate from preventive care visits. Additional visits also may become necessary if circumstances suggest concerns.

These recommendations represent a consensus by the American Academy of Pediatrics (AAP) and Bright Futures. The AAP continues to emphasize the great importance of continuity of care in comprehensive health supervision and the need to avoid fragmentation of care.

Refer to the specific guidance by age as listed in the *Bright Futures Guidelines* (Hagan JF, Shaw JS, Duncan PM, eds. *Bright Futures: Guidelines for Health Supervision of Infants, Children, and Adolescents*. 4th ed. American Academy of Pediatrics; 2017).

The recommendations in this statement do not indicate an exclusive course of treatment or serve as a standard of medical care. Variations, taking into account individual circumstances, may be appropriate.

The Bright Futures/American Academy of Pediatrics Recommendations for Preventive Pediatric Health Care are updated annually.

Copyright © 2024 by the American Academy of Pediatrics, updated June 2024.

No part of this statement may be reproduced in any form or by any means without prior written permission from the American Academy of Pediatrics except for one copy for personal use.

AGE ¹	INFANCY									EARLY CHILDHOOD							MIDDLE CHILDHOOD					ADOLESCENCE												
	Prenatal ¹	Newborn ³	3-5 d ⁴	By 1 mo	2 mo	4 mo	6 mo	9 mo	12 mo	15 mo	18 mo	24 mo	30 mo	3 y	4 y	5 y	6 y	7 y	8 y	9 y	10 y	11 y	12 y	13 y	14 y	15 y	16 y	17 y	18 y	19 y	20 y	21 y		
HISTORY	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	
MEASUREMENTS																																		
Length/Height and Weight		●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	
Head Circumference		●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	
Weight for Length		●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	
Body Mass Index ⁵		●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	
Blood Pressure ⁶		★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	
SENSORY SCREENING																																		
Vision ⁷		★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★
Hearing		● ⁸	● ⁹	→	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	
DEVELOPMENTAL/SOCIAL/BEHAVIORAL/MENTAL HEALTH																																		
Maternal Depression Screening ¹¹				●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●
Developmental Screening ¹²								●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●
Autism Spectrum Disorder Screening ¹³										●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●
Developmental Surveillance		●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●
Behavioral/Social/Emotional Screening ¹⁴		●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●
Tobacco, Alcohol, or Drug Use Assessment ¹⁵																						★	★	★	★	★	★	★	★	★	★	★	★	
Depression and Suicide Risk Screening ¹⁶																						●	●	●	●	●	●	●	●	●	●	●	●	
PHYSICAL EXAMINATION ¹⁷		●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●
PROCEDURES ¹⁸																																		
Newborn Blood		● ¹⁹	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	
Newborn Bilirubin ²⁰		●	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	
Critical Congenital Heart Defect ²¹		●	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	
Immunization ²²		●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●
Anemia ²⁴						★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★
Lead ²⁵						★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★
Tuberculosis ²⁷				★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★
Dyslipidemia ²⁸																						★	★	★	★	★	★	★	★	★	★	★	★	
Sexually Transmitted Infections ²⁹																						★	★	★	★	★	★	★	★	★	★	★	★	
HIV ³⁰																						★	★	★	★	★	★	★	★	★	★	★	★	
Hepatitis B Virus Infection ³¹		★	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	
Hepatitis C Virus Infection ³²																																		
Sudden Cardiac Arrest/Death ³³																						★	★	★	★	★	★	★	★	★	★	★	★	
Cervical Dysplasia ³⁴																																		
ORAL HEALTH ³⁵							● ³⁶	● ³⁶	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	
Fluoride Varnish ³⁷							→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	→	
Fluoride Supplementation ³⁸							★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★
ANTICIPATORY GUIDANCE	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	

1. If a child comes under care for the first time at any point on the schedule, or if any items are not accomplished at the suggested age, the schedule should be brought up to date at the earliest possible time.
2. A prenatal visit is recommended for parents who are at high risk, for first-time parents, and for those who request a conference. The prenatal visit should include anticipatory guidance, pertinent medical history, and a discussion of benefits of breastfeeding and planned method of feeding, per "The Prenatal Visit" (<https://doi.org/10.1542/peds.2019-1219>).
3. Newborns should have an evaluation after birth, and breastfeeding should be encouraged (and instruction and support should be offered).
4. Newborns should have an evaluation within 3 to 5 days of birth and within 48 to 72 hours after discharge from the hospital to include evaluation for feeding and jaundice. Breastfeeding newborns should receive formal breastfeeding evaluation, and their mothers should receive encouragement and instruction, as recommended in "Policy Statement: Breastfeeding and the Use of Human Milk" (<https://doi.org/10.1542/peds.2022-057988>). Newborns discharged less than 48 hours after delivery must be examined within

48 hours of discharge, per "Hospital Stay for Healthy Term Newborn Infants" (<https://doi.org/10.1542/peds.2015-0699>).
5. Screen, per "Clinical Practice Guideline for the Evaluation and Treatment of Children and Adolescents with Obesity" (<https://doi.org/10.1542/peds.2022-060640>).
6. Screening should occur per "Clinical Practice Guideline for Screening and Management of High Blood Pressure in Children and Adolescents" (<https://doi.org/10.1542/peds.2017-1909>). Blood pressure measurement in infants and children with specific risk conditions should be performed at visits before age 3 years.
7. A visual acuity screen is recommended at ages 4 and 5 years, as well as in cooperative 3-year-olds. Instrument-based screening may be used to assess risk at ages 12 and 24 months, in addition to the well visits at 3 through 5 years of age. See "Visual System Assessment in Infants, Children, and Young Adults by Pediatricians" (<https://doi.org/10.1542/peds.2015-3597>) and "Procedures for the Evaluation of the Visual System by Pediatricians" (<https://doi.org/10.1542/peds.2015-3597>).
8. Confirm initial screen was completed, verify results, and follow up, as appropriate. Newborns should be screened, per "Year 2007 Position Statement: Principles and Guidelines for Early Hearing Detection and Intervention Programs" (<https://doi.org/10.1542/peds.2007-2333>).

9. Verify results as soon as possible, and follow up, as appropriate.
10. Screen with audiometry including 6,000 and 8,000 Hz high frequencies once between 11 and 14 years, once between 15 and 17 years, and once between 18 and 21 years. See "The Sensitivity of Adolescent Hearing Screens Significantly Improves by Adding High Frequencies" (<https://www.sciencedirect.com/science/article/abs/pii/S1054139X16000483>).
11. Screening should occur per "Incorporating Recognition and Management of Perinatal Depression Into Pediatric Practice" (<https://doi.org/10.1542/peds.2018-3259>).
12. Screening should occur per "Promoting Optimal Development: Identifying Infants and Young Children With Developmental Disorders Through Developmental Surveillance and Screening" (<https://doi.org/10.1542/peds.2019-3449>).
13. Screening should occur per "Identification, Evaluation, and Management of Children With Autism Spectrum Disorder" (<https://doi.org/10.1542/peds.2019-3447>).

KEY: ● = to be performed ★ = risk assessment to be performed with appropriate action to follow, if positive ← ● or ★ → = range during which a service may be provided

(continued)

(continued)

14. Screen for behavioral and social-emotional problems per "Promoting Optimal Development: Screening for Behavioral and Emotional Problems" (<https://doi.org/10.1542/peds.2014-3710>), "Mental Health Competencies for Pediatric Practice" (<https://doi.org/10.1542/peds.2019-2757>), "Clinical Practice Guideline for the Assessment and Treatment of Children and Adolescents With Anxiety Disorders" (<https://pubmed.ncbi.nlm.nih.gov/32439401/>), "Screening for Anxiety in Adolescent and Adult Women: A Recommendation From the Women's Preventive Services Initiative" (<https://pubmed.ncbi.nlm.nih.gov/32510990/>), and "Anxiety in Children and Adolescents: Screening" (<https://www.uspreventiveservicestaskforce.org/uspstf/recommendation/screening-anxiety-children-adolescent>). The screening should be family centered and may include asking about caregiver emotional and mental health concerns and social determinants of health, racism, poverty, and relational health. See "Poverty and Child Health in the United States" (<https://doi.org/10.1542/peds.2016-0939>), "The Impact of Racism on Child and Adolescent Health" (<https://doi.org/10.1542/peds.2019-1765>), and "Preventing Childhood Toxic Stress: Partnering With Families and Communities to Promote Relational Health" (<https://doi.org/10.1542/peds.2021-052582>).
15. A recommended tool to assess use of alcohol, tobacco and nicotine, marijuana, and other substances, including opioids is available at <http://craftit.org>. If there is a concern for substance or opioid use, providers should consider recommending or prescribing Naloxone (see <https://www.cdc.gov/ore/search/pages/2018-evidence-based-strategies.html>) and <https://nida.nih.gov/publications/factsheets/naloxone>).
16. Screen adolescents for depression and suicide risk, making every effort to preserve confidentiality of the adolescent. See "Guidelines for Adolescent Depression in Primary Care (GLAD-PC): Part I. Practice Preparation, Identification, Assessment, and Initial Management" (<https://doi.org/10.1542/peds.2017-4081>), "Mental Health Competencies for Pediatric Practice" (<https://doi.org/10.1542/peds.2019-2757>), "Suicide and Suicide Attempts in Adolescents" (<https://doi.org/10.1542/peds.2016-1420>), and "The 21st Century Cures Act & Adolescent Confidentiality" (https://adolescenthealth.org/press_release/naspaq-sahm-statement-the-21st-century-cures-act-adolescent-confidentiality).
17. At each visit, age-appropriate physical examination is essential, with infant totally undressed and older children undressed and suitably draped. See "Use of Chaperones During the Physical Examination of the Pediatric Patient" (<https://doi.org/10.1542/peds.2011-0322>).
18. These may be modified, depending on entry point into schedule and individual need.
19. Confirm initial screen was accomplished, verify results, and follow up, as appropriate. The Recommended Uniform Screening Panel (<https://www.hrsa.gov/advisory-committees/heritable-disorders/usp/index.html>), as determined by The Secretary's Advisory Committee on Heritable Disorders in Newborns and Children, and state newborn screening laws/regulations (<https://www.babyfirsttest.org/>) establish the criteria for and coverage of newborn screening procedures and programs.
20. Verify results as soon as possible, and follow up, as appropriate.
21. Confirm initial screen was accomplished, verify results, and follow up, as appropriate. See "Clinical Practice Guideline Revision: Management of Hyperbilirubinemia in the Newborn Infant 35 or More Weeks of Gestation" (<https://doi.org/10.1542/peds.2022-058859>).
22. Screening for critical congenital heart disease using pulse oximetry should be performed in newborns, after 24 hours of age, before discharge from the hospital, per "Endorsement of Health and Human Services Recommendation for Pulse Oximetry Screening for Critical Congenital Heart Disease" (<https://doi.org/10.1542/peds.2011-3211>).
23. Schedules, per the AAP Committee on Infectious Diseases, are available at <https://publications.aap.org/redbook/pages/immunization-schedules>. Every visit should be an opportunity to update and complete a child's immunizations.
24. Perform risk assessment or screening, as appropriate, per recommendations in the current edition of the AAP *Pediatric Nutrition: Policy of the American Academy of Pediatrics* (Iron chapter).
25. For children at risk of lead exposure, see "Prevention of Childhood Lead Toxicity" (<https://doi.org/10.1542/peds.2016-1493>) and "Low Level Lead Exposure Harms Children: A Renewed Call for Primary Prevention" (<https://stacks.cdc.gov/view/cdc/11859>).

26. Perform risk assessments or screenings as appropriate, based on universal screening requirements for patients with Medicaid or in high prevalence areas.
27. Tuberculosis testing per recommendations of the AAP Committee on Infectious Diseases, published in the current edition of the AAP *Red Book: Report of the Committee on Infectious Diseases*. Testing should be performed on recognition of high-risk factors.
28. See "Integrated Guidelines for Cardiovascular Health and Risk Reduction in Children and Adolescents" (<http://www.nhlbi.nih.gov/guidelines/cvd/index.html>).
29. Adolescents should be screened for sexually transmitted infections (STIs) per recommendations in the current edition of the AAP *Red Book: Report of the Committee on Infectious Diseases*.
30. Screen adolescents for HIV at least once between the ages of 15 and 21, making every effort to preserve confidentiality of the adolescent, as per "Human Immunodeficiency Virus (HIV) Infection: Screening" (<https://www.uspreventiveservicestaskforce.org/uspstf/recommendation/human-immunodeficiency-virus-hiv-infection-screening>); after initial screening, youth at increased risk of HIV infection should be retested annually or more frequently, as per "Adolescents and Young Adults: The Pediatrician's Role in HIV Testing and Pre- and Postexposure HIV Prophylaxis" (<https://doi.org/10.1542/peds.2021-055207>).
31. Perform a risk assessment for hepatitis B virus (HBV) infection according to recommendations per the USPSTF (<https://www.uspreventiveservicestaskforce.org/uspstf/recommendation/hepatitis-b-virus-infection-screening>) and in the 2021-2024 edition of the AAP *Red Book: Report of the Committee on Infectious Diseases*, making every effort to preserve confidentiality of the patient.
32. All individuals should be screened for hepatitis C virus (HCV) infection according to the USPSTF (<https://www.uspreventiveservicestaskforce.org/uspstf/recommendation/hepatitis-c-screening>) and Centers for Disease Control and Prevention (CDC) recommendations (<https://www.cdc.gov/mmwr/volumes/69/rr/r6902a.htm>) at least once between the ages of 18 and 79. Those at increased risk of HCV infection, including those who are persons with past or current injection drug use, should be tested for HCV infection and reassessed annually.
33. Perform a risk assessment, as appropriate, per "Sudden Death in the Young: Information for the Primary Care Provider" (<https://doi.org/10.1542/peds.2021-052049>).
34. See USPSTF recommendations (<https://www.uspreventiveservicestaskforce.org/uspstf/recommendation/cervical-cancer-screening>). Indications for pelvic examinations prior to age 21 are noted in "Gynecologic Examination for Adolescents in the Pediatric Office Setting" (<https://doi.org/10.1542/peds.2010-1564>).
35. Assess whether the child has a dental home. If no dental home is identified, perform a risk assessment (<https://www.aap.org/en/patient-care/oral-health/oral-health-practice-tools>) and refer to a dental home. Recommend brushing with fluoride toothpaste in the proper dosage for age. See "Maintaining and Improving the Oral Health of Young Children" (<https://doi.org/10.1542/peds.2022-060417>).
36. Perform a risk assessment (<https://www.aap.org/en/patient-care/oral-health/oral-health-practice-tools>). See "Maintaining and Improving the Oral Health of Young Children" (<https://doi.org/10.1542/peds.2022-060417>).
37. The USPSTF recommends that primary care clinicians apply fluoride varnish to the primary teeth of all infants and children starting at the age of primary tooth eruption (<https://www.uspreventiveservicestaskforce.org/uspstf/recommendation/prevention-of-dental-caries-in-children-younger-than-age-5-years-screening-and-interventions>). Once teeth are present, apply fluoride varnish to all children every 3 to 6 months in the primary care or dental office based on caries risk. Indications for fluoride use are noted in "Fluoride Use in Caries Prevention in the Primary Care Setting" (<https://doi.org/10.1542/peds.2020-034637>).
38. If primary water source is deficient in fluoride, consider oral fluoride supplementation. See "Fluoride Use in Caries Prevention in the Primary Care Setting" (<https://doi.org/10.1542/peds.2020-034637>).

Summary of Changes Made to the Bright Futures/AAP Recommendations for Preventive Pediatric Health Care (Periodicity Schedule)

This schedule reflects changes approved in December 2023 and published in June 2024. For updates and a list of previous changes made, visit www.aap.org/periodicityschedule.

FOOTNOTE CHANGES MADE IN DECEMBER 2023

- **3-5 DAY VISIT (Footnote 4)**
This footnote reflects the AAP "Policy Statement: Breastfeeding and the Use of Human Milk", published June 2022.
- **BODY MASS INDEX (Footnote 5)**
This footnote reflects the AAP "Clinical Practice Guideline for the Evaluation and Treatment of Children and Adolescents with Obesity", published January 2023.
- **BEHAVIORAL/SOCIAL/EMOTIONAL SCREENING (Footnote 14)**
This footnote reflects the USPSTF "Anxiety in Children and Adolescents: Screening" recommendations, published October 2022.
- **TOBACCO, ALCOHOL, OR DRUG USE ASSESSMENT (Footnote 15)**
This footnote reflects the Centers for Disease Control (CDC) and National Institute of Drug Abuse (NIDA) guidance related to recommending and prescribing Naloxone.
- **NEWBORN BILIRUBIN SCREENING (Footnote 21)**
This footnote reflects the AAP "Clinical Practice Guideline Revision: Management of Hyperbilirubinemia in the Newborn Infant 35 or More Weeks of Gestation", published August 2022.
- **ORAL HEALTH (Footnotes 35 and 36)**
These footnotes reflect the AAP clinical report, "Maintaining and Improving the Oral Health of Young Children", published December 2022.

CHANGES MADE IN DECEMBER 2022

HIV

The HIV screening recommendation has been updated to extend the upper age limit from 18 to 21 years (to account for the range in which the screening can take place) to align with recommendations of the US Preventive Services Task Force and AAP policy ("Adolescents and Young Adults: The Pediatrician's Role in HIV Testing and Pre- and Postexposure HIV Prophylaxis").

- Footnote 30 has been updated to read as follows: "Screen adolescents for HIV at least once between the ages of 15 and 21, making every effort to preserve confidentiality of the adolescent, as per "Human Immunodeficiency Virus (HIV) Infection: Screening" (<https://www.uspreventiveservicestaskforce.org/uspstf/recommendation/human-immunodeficiency-virus-hiv-infection-screening>); after initial screening, youth at increased risk of HIV infection should be retested annually or more frequently, as per "Adolescents and Young Adults: The Pediatrician's Role in HIV Testing and Pre- and Postexposure HIV Prophylaxis" (<https://doi.org/10.1542/peds.2021-055207>)"



HRSA
www.hrsa.gov

This program is supported by the Health Resources and Services Administration (HRSA) of the U.S. Department of Health and Human Services (HHS) as part of an award totaling \$1,766,600 with 0% financed with non-governmental sources. The contents of this document are those of the author(s) and do not necessarily represent the official views of, nor an endorsement by, HRSA, HHS, or the U.S. Government. For more information, please visit HRSA.gov.